

Relativistic unitary coupled cluster theory and applications

Chiranjib Sur, Rajat K Chaudhuri, Bijaya K. Sahoo, B. P. Das
Non-Accelerator Particle Physics Group, Indian Institute of Astrophysics, Bangalore - 560 034, India

D. Mukherjee
Department of Physical Chemistry, Indian Association for the Cultivation of Science, Kolkata - 700 032, India

(Dated: December 14, 2004)

We present the first formulation and application of relativistic unitary coupled cluster theory to atomic properties. The remarkable features of this theory are highlighted, and it is used to calculate the lifetimes of $5^2D_{3/2}$ and $6^2P_{3/2}$ states of Ba^+ and Pb^+ respectively. The results clearly suggest that it is very well suited for accurate *ab initio* calculations of properties of heavy atomic systems.

PACS numbers: 31.15.Ar, 31.15.Dv, 31.25.Jf, 32.10.Fn

There have been a number of attempts to modify coupled-cluster (CC) theory [1], despite its spectacular success in elucidating the properties of a wide range of many-body systems [1, 2, 3, 4, 5]. One interesting case in point is unitary coupled-cluster (UCC) theory which was first proposed by Kutzelnigg [6]. In this theory, the effective Hamiltonian is Hermitian by construction and the energy which is the expectation value of this operator in the reference state is unlike in CC theory, an upper bound to the ground state energy [5]. Another attractive feature of this theory which we shall discuss later is that at a given level of approximation it incorporates certain higher order excitations that are not present in CC theory. Furthermore, it is well suited for the calculation of properties where core relaxation effects are important [7, 8]. In spite of the aforementioned advantages, there have been relatively few studies based on this method [9, 10]. This work is the first relativistic formulation of unitary coupled cluster (UCC) theory and also the first application of this theory to atomic properties. In this letter, we first present the formal aspects of relativistic UCC theory and then apply it to calculate the lifetimes of the $5^2D_{3/2}$ and $6^2P_{3/2}$ states of Ba^+ and Pb^+ respectively; which depend strongly on both relativistic and correlation effects. The comparison of the results of these calculations with accurate experimental data would constitute an important test of this theory.

The exact wave function for a closed shell state in CC theory is obtained by the action of the operator $\exp(T)$ on the reference state $|\Phi\rangle$. However, in UCC theory [10], it is written as

$$|\Psi\rangle = \exp(\sigma) |\Phi\rangle \quad (1)$$

where $\sigma = T - T^\dagger$; and T and $T^\dagger$ are the excitation and deexcitation operators respectively. σ is clearly anti-Hermitian, since $\sigma^\dagger = -\sigma$. Using this unitary ansatz for the correlated wave function, the relativistic UCC equation

in the Dirac-Coulomb approximation can be written as

$$\exp(\sigma^\dagger) H \exp(\sigma) |\Phi\rangle = E |\Phi\rangle, \quad (2)$$

where H is the Dirac-Coulomb Hamiltonian

$$H = \sum_i c\alpha_i \cdot p_i + (\beta - 1)mc^2 + V_N + \sum_{i<j} \frac{e^2}{r_{ij}}. \quad (3)$$

Using the normal ordered Hamiltonian, Eq.(2) can be rewritten as

$$\exp(\sigma^\dagger) H_N \exp(\sigma) |\Phi\rangle = \Delta E |\Phi\rangle, \quad (4)$$

where the normal ordered Hamiltonian is defined as $H_N = H - \langle\Phi|H|\Phi\rangle$ and $\Delta E = E - \langle\Phi|H|\Phi\rangle$.

The choice of the operator σ makes the effective Hamiltonian $\bar{H}_N = \exp(-\sigma) H_N \exp(\sigma)$ Hermitian.

The effective Hamiltonian is expressed by the Hausdorff expansion in CC theory as

$$\begin{aligned} \bar{H}_N &= \exp(-T) H_N \exp(T) \\ &= H_N + [H_N, T] + \frac{1}{2!} [[H_N, T], T] \\ &\quad + \frac{1}{3!} [[[H_N, T], T], T] \\ &\quad + \frac{1}{4!} [[[[H_N, T], T], T], T]. \end{aligned} \quad (5)$$

In UCC, the operator T is replaced by $\sigma = T - T^\dagger$ in the above equation and this results in $\bar{H}_N$ being expressed in terms of a non-terminating series of commutators. For practical reasons, one truncates the series after some finite order. Truncation at the n -th order commutator leads to the nomenclature UCC(n).

Using UCC(3) approximation and without modifying the last term of Eq. (5), we write Eq. (4) as

$$\left[H_N + \overline{H_N T} + \frac{1}{2!} \left(\overline{H_N T T} + 2 \overline{T^\dagger H_N T} \right) + \frac{1}{3!} \left(\overline{H_N T T T} + 3 \overline{T^\dagger T^\dagger H_N T} + 3 \overline{T^\dagger H_N T T} \right) + \frac{1}{4!} \overline{H_N T T T T} \right] |\Phi\rangle = \Delta E |\Phi\rangle \quad (6)$$

By projecting single, double and higher order excited determinant states on Eq.(6), we get the cluster amplitude equations. The approximation which includes only single and double excitations/deexcitations in the UCC wavefunction is known as UCCSD. After careful analysis, one finds that there are new terms arising from $\overline{T^\dagger H_N T}$, $\overline{T^\dagger T^\dagger H_N T}$ and $\overline{T^\dagger H_N T T}$ in the UCCSD equations. The first term will give rise to some extra diagrams which correspond to double and triple excitations and the last two terms to triple and quadrupole excitations. Some typical diagrams which represent triple and quadrupole excitations are given in figures 1a and 1b respectively. The UCCSD approximation in addition to single and double excitations, also includes some triple and quadrupole excitations to all orders in the residual Coulomb interaction in a more elegant and simpler manner compared to CC theory. It would have been computationally prohibitive to include triple and quadrupole excitations for heavy atoms in the framework of normal CC theory. Indeed one of the principal advantages of UCC theory is its ability to subsume higher levels of excitations than CC theory at the same level of approximation.

To calculate the ground state of the system we first compute the cluster amplitude of the closed shell states (Ba^{++} and Pb^{++}) by using Eq.(6) and then use the open shell coupled cluster method (OSCC)[11]. The exact wave function can then be written as

$$|\Psi_k^{N+1}\rangle = \exp(\sigma) \{1 + S_k\} |\Phi_k^{N+1}\rangle, \quad (7)$$

where $|\Phi_k^{N+1}\rangle$ is the Dirac-Fock reference state which we get after adding an electron to the k th virtual orbital and S_k is the corresponding excitation operator. We obtain a set of equations

$$\langle \Phi_k^{N+1} | \overline{H_N} (1 + S_k) | \Phi_k^{N+1} \rangle = H_{\text{eff}} \quad (8)$$

and

$$\begin{aligned} & \langle \Phi_k^{*N+1} | \overline{H_N} (1 + S_k) | \Phi_k^{N+1} \rangle \\ &= \langle \Phi_k^{*N+1} | S_k | \Phi_k^{N+1} \rangle H_{\text{eff}}. \end{aligned} \quad (9)$$

The Eq.(9) is non-linear in S_k because H_{eff} is itself a function of S_k where $S_k = \sum_{ak} s_a^k a_k^\dagger a_a + \sum_{abkr} s_{ab}^{kr} a_k^\dagger a_r^\dagger a_b a_a$; s_a^k and s_{ab}^{kr} are the single and double excitation cluster amplitudes for the valence electrons. The labels a, b and k, r refer to the core and virtual orbitals respectively. Hence, these equations have to be solved self-consistently to determine the S_k amplitudes. We have included the triple excitations in our calculations in an approximate

way. The amplitudes corresponding to these excitations are of the form

$$S_{abk}^{pqr} = \frac{\widehat{VT}_2 + \widehat{VS}_2}{\varepsilon_a + \varepsilon_b + \varepsilon_k - \varepsilon_p - \varepsilon_q - \varepsilon_r}, \quad (10)$$

where S_{abk}^{pqr} are the amplitudes corresponding to the simultaneous excitation of orbitals a, b, k to p, q, r respectively and $\widehat{VT}$ and $\widehat{VS}$ are the correlated composites involving V and T , and V and S respectively. ε 's are the orbital energies. The above amplitudes are added appropriately in the singles and doubles open shell cluster amplitude equations and they are then solved self-consistently. We therefore obtain solutions of S_1 and S_2 amplitudes taking into consideration the partial effect of the triple excitations. We had referred to this approximation earlier as CCSD(T).

As we have seen in Eq.(6), $\overline{H_N}$ is defined as $\overline{H_N} = \exp(-\sigma) H_N \exp(\sigma)$. From Eq.(5) it is clear that the expansion of $\overline{H_N}$ corresponding to UCC theory can be expressed as

$$\overline{H_N} = (\overline{H_N})_{\text{CCSD}} + (\overline{H_N})_{\text{extra}}. \quad (11)$$

Following much the same argument as in the closed shell case, it can be shown that the contributions from the extra part arising in Eq.(11) contains terms like $\overline{T^\dagger H_N T}$, $\overline{T^\dagger T^\dagger H_N T}$ and $\overline{T^\dagger H_N T T}$. These terms will affect the open-shell amplitude determining equations. In the UCC theory for the closed shell, we have seen that although there are some extra terms arising from the expansion of $\overline{H_N}$, the contributing diagrams corresponding to those terms are not always new. Hence we have taken into account those diagrams which are not present in the CCSD approximation. It can be shown that the extra terms arising from the inclusion on the operator $\sigma = T - T^\dagger$ give rise to some new terms but the corresponding diagrams have already been taken into account through the inclusion of partial triple amplitudes. In analogy with CCSD(T), we refer to our approach as UCCSD(T).

The normalized transition matrix element ($i \rightarrow f$) due to an operator $\hat{O}$ is given by

$$\hat{O}_{fi} = \frac{\langle \Psi_f^{N+1} | \hat{O} | \Psi_i^{N+1} \rangle}{\sqrt{\langle \Psi_f^{N+1} | \Psi_f^{N+1} \rangle \langle \Psi_i^{N+1} | \Psi_i^{N+1} \rangle}}$$

We consider next the calculations of the lifetimes of the $5^2D_{3/2}$ and $6^2P_{3/2}$ states of Ba^+ and Pb^+ respectively.





FIG. 1: Typical triples quadruples diagrams arising from UCC(3)

FIG. 2: Decay scheme of the low lying states of Ba^+ and Pb^+

Both of these states decay to the ground states via $M1$ and $E2$ transitions as shown in figure 2. The transition probabilities A ($A_{fi} = A_{i \rightarrow f}$ in s^{-1}) for $M1$ and $E2$ transitions are expressed as [12]

$$A_{fi}^{M1} = \frac{2.6973 \times 10^{13}}{[J_i] \lambda^3} S_{fi}^{M1} \quad (12)$$

and

$$A_{fi}^{E2} = \frac{1.1199 \times 10^{18}}{[J_i] \lambda^5} S_{fi}^{E2} \quad (13)$$

respectively, where J_i is the degeneracy of the initial state and λ (in $\text{\AA}$) is the wavelength corresponding to the transition $i \rightarrow f$. In Eqs.(12) and (13) the line-strength S_{fi} is $|M1_{fi}|^2$ and $|E2_{fi}|^2$ respectively and $M1_{fi}$ and $E2_{fi}$ are the corresponding one electron reduced matrix elements of magnetic dipole and electric quadrupole transitions [13].

The net probability for a given transition which allows two different channels is given by [12]

$$A = A_{M1} + A_{E2} \quad (14)$$

and the corresponding lifetime, which is the inverse of the transition probability is expressed as

$$\frac{1}{\tau} = \frac{1}{\tau_{M1}} + \frac{1}{\tau_{E2}}. \quad (15)$$

The results of the calculations of the lifetimes of the $5^2D_{3/2}$ state of Ba^+ and $6^2P_{3/2}$ state of Pb^+ are given in tables II and III respectively. These calculations are

TABLE I: Description of the basis functions used in the UCC calculation of Ba^+ .

	$s_{1/2}$	$p_{1/2}$	$p_{3/2}$	$d_{3/2}$	$d_{5/2}$	$f_{5/2}$	$f_{7/2}$	$g_{7/2}$	$g_{9/2}$
Analytical	5	6	6	5	5	8	8	6	6
Numerical	8	7	7	5	5	2	2	0	0
Total	13	13	13	10	10	10	10	6	6

TABLE II: Excitation energies (in cm^{-1}) and lifetime (in s) of $5^2D_{3/2}$ state of Ba^+ corresponding to the transition $5^2D_{3/2} \rightarrow 6^2S_{1/2}$

	Excitation energy	τ
UCCSD(T)	4789	81.01
CCSD(T) [14]	4809	87.06
Dzuba <i>et al</i> [15]	4411	81.5
Guet <i>et al</i> [16]	4688	83.7
Experiment [17]	4874	79 ± 4.6

very challenging as they involve accurate determinations of the line strengths of the $M1$ and $E2$ transitions and more critically the third and fifth powers of the excitation energies.

The lifetime of the $5^2D_{3/2}$ state of Ba^+ is calculated using hybrid orbitals consisting of single particle orbitals which are partly numerical and partly analytical. Such an approach is described in detail by Majumder *et al* [18]. The analytical orbitals are Gaussian type orbitals (GTOs). The details of this basis are given in table I.

TABLE III: Excitation energies (in cm^{-1}) and lifetime (in s) of the $6^2P_{3/2}$ state of Pb^+

	Excitation energy	τ
Dirac-Fock	13612	0.0415
CCSD(T)[13]	13710	0.0425
UCCSD(T)	13719	0.0413
Exp[20]	14085	.0412(7)

Excitations from all the core orbitals are included for this ion as well as Pb^+ . Since the calculation of transition probabilities/lifetimes are extremely sensitive to the excitation energies. Gopakumar *et al* [14] have shown that the hybrid basis can serve this purpose very well. Our UCCSD(T) calculations yield ionization potentials (IPs) to an accuracy of about 0.1%, and 0.07% for the $6^2S_{1/2}$ and $5^2D_{3/2}$ states of Ba^+ respectively. The corresponding IP (in cm^{-1}) values are 80544 (80687) and 75755 (75813). The numbers in the parentheses are the corresponding experimental values [19].

The values of the calculated and experimental lifetimes are given in table II. Dzuba *et al*[15], Guet *et al* [16] and Gopakumar *et al* [14] have not considered the $M1$ channel in their calculations. Gopakumar *et al* have used CCSD(T), but Guet *et al* and Dzuba *et al* have used different variants of many-body perturbation theory (MBPT) with certain semi-empirical features to calculate the transition amplitudes. In addition, the latter two have used experimental values of the excitation energies to calculate the transition probabilities/lifetimes. Our calculation based on UCCSD(T) is purely *ab initio* and takes into account the contribution of the $M1$ transition. The transition amplitude in this approach includes some terms in addition to those that appear in

CCSD(T); and they arise due to the presence of the deexcitation operator in the UCC wavefunction. The result of the lifetime calculation considering only the $E2$ channel is 85.567. Inclusion of the $M1$ transition improves the total value (81.01 s) which is within the experimental error bar (79 ± 4.6 s).

The UCCSD(T) calculations of lifetime for the $6^2P_{3/2}$ state of Pb^+ are carried out using the same basis functions used by Sahoo *et al* [13]. The leading correlation contribution to the $M1$ and $E2$ channel come from the core polarization and pair correlation effects. It is evident from Table III that the accuracy of the calculation of excitation energies using UCCSD(T) is even better than that obtained from CCSD(T). This result and the improvements in the $M1$ and $E2$ transition amplitudes due to the UCC formulation coupled together give a value of the lifetime of the $6^2P_{3/2}$ state of Pb^+ that is clearly in better agreement with the corresponding CCSD(T) calculation. This value is within the limits of the experimental error [20], but this is not the case for CCSD(T).

In summary, we have applied the relativistic UCC theory for the first time to atomic properties. The results of our UCCSD(T) calculations of the lifetimes of the $5^2D_{3/2}$ and $6^2P_{3/2}$ states of Ba^+ and Pb^+ respectively are in very good agreement with experiment and superior to those of all previous calculations. It indeed appears that this theory is capable of yielding high precision results for a wide range of properties of heavy atomic systems including violation of parity and time-reversal symmetries. In addition, it would be worthwhile to explore its feasibility for the accurate determination of the properties of other many-body systems, particularly when it becomes necessary to go beyond the usual coupled cluster theory.

This work was supported by the BRNS for project no. 2002/37/12/BRNS. The computations were done on our group's Xeon PC cluster and Param Padma, the Teraflop Supercomputer in CDAC, Bangalore. We thank Professor Güenther Werth for useful discussions.

-
- | | |
|--|---|
| <p>[1] R. F. Bishop, <i>Microscopic Quantum Many-Body Theories and Their Applications</i>, p.1, Ed. J. Navarro and A. Polls, Springer Verlag, Berlin, (1998).</p> <p>[2] I. Lindgren and J. Morrison, <i>Atomic Many-Body Theory</i>, Springer, Berlin (1985).</p> <p>[3] U. Kaldor, <i>Microscopic Quantum Many-Body Theories and Their Applications</i>, p.71, Ed. J. Navarro and A. Polls, Springer Verlag, Berlin, (1998).</p> <p>[4] R. J. Bartlett, <i>Modern Electronic Structure Theory</i>, vol-II, p.1047, Ed. D. R. Yarkony, World Scientific, Singapore (1995).</p> <p>[5] T. D. Crawford and H. F. Schaefer III, <i>Rev. Comput. Chem.</i>, 14, 33 (2000).</p> <p>[6] W. Kutzelnigg, <i>J. Chem. Phys.</i>, 77, 3081 (1982).</p> <p>[7] K. P. Geetha, <i>Ph. D Thesis</i>, Bangalore University (2001).</p> <p>[8] C. Sur and R. K. Chaudhuri, <i>J. Phys. B</i>, 37, 4127 (2004).</p> <p>[9] J. D. Watts, G. W. Trucks and R. J. Bartlett, <i>Chem.</i></p> | <p><i>Phys. Lett.</i>, 157, 359 (1989).</p> <p>[10] M. R. Hoffmann and J. Simons, <i>J. Chem. Phys.</i>, 88, 993 (1988).</p> <p>[11] C. Sur, B. K. Sahoo, R. K. Chaudhuri, B. P. Das and D. Mukherjee, To be published in <i>Eur. Phys. J. D</i> (2004) and the references therein.</p> <p>[12] W. C. Martin and W. L. Wiese, p. 135, <i>Atomic, Molecular and Optical Physics Handbook</i>, ed. W. F. Drake Gordon, AIP Press, Woodburg, NY (1996).</p> <p>[13] B. K. Sahoo, S. Majumder, R. K. Chaudhuri, B. P. Das and D. Mukherjee, <i>J. Phys. B</i>, 37, 3409 (2004).</p> <p>[14] G. Gopakumar, H. Merlitz, R. K. Chaudhuri, B. P. Das, U. S. Mahapatra and D. Mukherjee, <i>Phys. Rev. A</i>, 66, 032505 (2002).</p> <p>[15] V. A. Dzuba, V. V. Flambaum and J. S. M. Ginges, <i>Phys. Rev. A</i>, 63, 062101 (2001).</p> <p>[16] C. Guet and W. R. Johnson, <i>Phys. Rev. A</i>, 44, 1531</p> |
|--|---|

- (1991).
- [17] N. Yu, W. Nagourney and H. Dehmelt, *Phys. Rev. Lett.*, **78**, 4898 (1997).
- [18] S. Majumder, G. Gopakumar, H. Merlitz and B. P. Das, *J. Phys. B*, **34**, 4821 (2001).
- [19] C. E. Moore, *Natl. Stand. Ref. Data Ser.* (U. S. Natl. Bur. Stand.), **3** (1971).
- [20] A. Roth, Ch Gerz, D. Wilsdorf and G. Werth, *Z. Phys. D*, **11**, 283 (1989).